Getting started with dualPortal Endoscopic Spine Surgery

James M. Mok, MD, MBA
Clinical Associate Professor, University of Illinois-Chicago
NorthShore University HealthSystem

2nd Annual Amplify Surgical Endoscopic Spine Symposium March 4, 2023









Unilateral Biportal Endoscopy









- Korean healthcare market
- Innovate to compete
- Implemented in US by Dr.
 Don Park






Primum Non Nocere

- N=7, 2-3+ hours
- No neuromonitoring changes (EMG/SSEP)
- Enormous amount of fluid (my record = 56 liters)
- Ensure good outflow to prevent extravasation into tissue
- RF probe used at Ablate=7 and Coag=2
- Diamond tip bur


Bur



Rated for water <50,000 RPM to minimize bone dust



Set-up





C-armor

Ergonomics

Good

- No microscope or loupes
- No tilting

Bad

- Carpal tunnel syndrome
- Arm fatigue




The first step is critical to determining success










Bleeding



- Increase water pressure
- Decrease SBP (anesthesia)
- RF probe
- Don't cause it in the first place







Getting started with dualPortal Endoscopic Spine Surgery



- 1. Well-established
- 2. Safe
- 3. Diamond tip bur <50,000 RPM
- 4. Set-up with Shoulder Pouch (2 drains)
- 5. Lower the table
- 6. Build a good fighting hole
- 7. Avoid bleeding in the first place
- 8. RF probe for bleeding
- 9. Your beacons: midline raphe and SAP



